

KNOW YOUR CIDP TREATMENT OPTIONS

IVIg

(intravenous immunoglobulin)

Frequency & Time*:

~Every 3 weeks for ~3-5 hours

Method:

IV infusion (administered by an HCP)

Location:

Hospital, doctor's office,
infusion center, home

SCIg

(subcutaneous immunoglobulin)

Frequency & Time*:

~Every week for ~1-1.5 hours

Method†:

SC infusion (administered by
an HCP or self-administered)

Location:

Hospital, doctor's office,
infusion center, home

STEROIDS

Frequency & Time*:

Daily or pulsed, and time depends
on individual

Method†:

Oral (self-administered) or
IV infusion (administered by an HCP)

Location:

Oral: Home

IV: Hospital, doctor's office,
infusion center, home

PLASMA EXCHANGE

Frequency & Time*:

Depends on individual

Method:

IV infusion (administered by an HCP)

Location:

Specialized infusion center

FcRn BLOCKERS

(neonatal Fc receptor blockers)

Frequency & Time*:

Every week for 20-90 seconds,
depending on administration method

Method†:

SC injection (administered by an HCP
or self-administered, depending on
treatment form)

Location: Doctor's office, home

*Frequency and time may vary and don't include preparation or monitoring time.

†All self-administered treatment options can also be administered by a caregiver.

HCP=healthcare professional.

EXPLORE A CIDP TREATMENT OPTION:

[ThisCIDPTreatment.com](https://www.thiscidptreatment.com)

*You'll be taken to a website to learn more
about this treatment.*

For U.S. audiences only.

©2026 argenx US-NON-26-00188 V1 09/2026

argenx 