

KNOW YOUR CIDP TREATMENT OPTIONS

Treatment	Frequency & Time*	Method†	Location
IVIg (intravenous immunoglobulin)	~Every 3 weeks for ~3-5 hours	IV infusion (administered by an HCP)	Hospital, doctor's office, infusion center, home
SCIg (subcutaneous immunoglobulin)	~Every week for ~1-1.5 hours	SC infusion (administered by an HCP or self-administered)	Hospital, doctor's office, infusion center, home
STEROIDS	Daily or pulsed, and time depends on individual	Oral (self-administered) or IV infusion (administered by an HCP)	Oral: Home IV: Hospital, doctor's office, infusion center, home
PLASMA EXCHANGE	Depends on individual	IV infusion (administered by an HCP)	Specialized infusion center
FcRn BLOCKERS (neonatal Fc receptor blockers)	Every week for 20-90 seconds, depending on administration method	SC injection (administered by an HCP or self-administered, depending on treatment form)	Doctor's office, home

*Frequency and time may vary and don't include preparation or monitoring time.

†All self-administered treatment options can also be administered by a caregiver.

HCP=healthcare professional.

EXPLORE A CIDP TREATMENT OPTION:

[ThisCIDPTreatment.com](https://www.thiscidptreatment.com)



*You'll be taken to a website to learn
more about this treatment.*